

HOW DERBY'S AIR POLLUTION IMPACTS OUR HEALTH AND PUTS A STRAIN ON THE NHS

New research conducted by the Environmental Research Group at King's College London has quantified the likely impacts of air pollution in Derby on a range of important health conditions, including heart attacks, hospitalisations for stroke and emergency asthma admissions amongst adults and children.

The research suggests that each year in Derby, higher air pollution days (compared to low pollution days) are linked to:

A 1.8% INCREASE IN THE RISK OF HAVING AN OUT OF HOSPITAL CARDIAC ARREST

AN EXTRA 5 CHILDREN BEING HOSPITALISED WITH ASTHMA

SENDING UP TO 13 MORE PEOPLE TO HOSPITAL FOR STROKE

THE REPORT ESTIMATES THAT, IN DERBY:

- Your child is 6.2% more likely to be hospitalised for asthma on days with high nitrogen dioxide pollution compared to days with lower air pollution

IF AIR POLLUTION IN DERBY WAS CUT BY JUST ONE FIFTH, EVERY YEAR WE WOULD SEE:

- 179 fewer children suffering with low lung function
- 3 fewer babies being born underweight
- Fewer children suffering with a chest infection and 85 fewer asthmatic children suffering with bronchitic symptoms (cough and phlegm)
- A decrease in the risk of coronary heart disease by around -% which would result in -fewer cases a year
- A decrease in lung cancer cases by around -%, which would result in -fewer cases every year

KEY POINTS

New research has estimated how the air pollution levels recorded in Derby impact on health in the city

The study estimates how current levels of pollution increase hospital admission rates, the risks of developing cancer, and worsen child health in the city

This study adds to the already compelling case for urgent and ambitious action to address air pollution in Derby

AIR POLLUTION IN DERBY

Air quality monitoring in Derby shows that the city is regularly above the EU and World Health Organisation guideline levels for Nitrogen Dioxide. To address this, an Air Quality Management Area has been declared. More information is available [here](#).

MEASURES WHICH CAN REDUCE AIR POLLUTION IN DERBY

There are a range of policies and interventions which can reduce air pollution in areas like Derby. These include measures such as:

- Promotion of active travel (walking and cycling), including through infrastructure improvements
- Electrification of vehicles, including both private vehicles and public ones such as bus fleets
- A move away from the use of private vehicles to public transport, alongside better traffic planning and management
- Shifts towards cleaner burning fuels and more modern engines
- Reductions in household solid fuel (including wood) burning.
- Addressing construction and industrial emissions through setting (and enforcing) higher standards.

For further information on air quality improvement interventions you can read the Public Health England Review of interventions [here](#). The government Clean Air Strategy 2019 is [here](#).

BACKGROUND

While there are a wealth of studies which demonstrate the links between poor air quality and adverse health outcomes, few studies have synthesised these for a number of health outcomes and 'at risk' groups like children. Fewer still have applied the emerging insight to cities across the UK drawing on the available monitoring data which is collected by DEFRA. This study is the first to attempt to do this and aims to localize the research so that policy makers and the public can be better informed about the nature of the local air pollution problem, to support local policies which can address this often invisible public health crisis.

This research was conducted by the Environmental Research Group at King's College London and was funded by [the Clean Air Fund](#).

LINKS TO THE CORE REPORT

Please click [here](#) for the core academic report on which this brief is based. Data from the UK Automatic Urban and Rural Network (AURN) are published by Defra [here](#).

METHODS

These statements 'personalise' the health effects of air pollution. They have been developed from calculations based on three components:

1. A measure of exposure: using air quality monitoring data from regulatory based monitoring networks.
2. A numerical relationship between the air pollutant concentration ('exposure') and the change in the health outcome in question. This numerical relationship is termed the 'concentration response function' or CRF, and has been drawn from a comprehensive review of air pollution research. It usually takes the form of a percentage increase in adverse health impacts over the baseline rate.
3. This percentage change in the health outcome due to pollutant exposure is applied to the baseline rate of the outcome or disease.

The result of this is used to develop quantitative statements giving the effect of a given exposure to an air pollutant on a particular health outcome or disease.

For a detailed description of methods, please consult the full King's College Report.

Produced by:



Drawing on research by:

